

ON THE DEGREES OF POLYNOMIAL DIVISORS OVER FINITE FIELDS

ANDREAS WEINGARTNER

ABSTRACT. We show that the proportion of polynomials of degree n over the finite field with q elements, which have a divisor of every degree below n , is given by $c_q n^{-1} + O(n^{-2})$. More generally, we give an asymptotic formula for the proportion of polynomials, whose set of degrees of divisors has no gaps of size greater than m . To that end, we first derive an improved estimate for the proportion of polynomials of degree n , all of whose non-constant divisors have degree greater than m . In the limit as $q \rightarrow \infty$, these results coincide with corresponding estimates related to the cycle structure of permutations.

1. INTRODUCTION

There are many parallels between the factorization of integers into primes and the decomposition of combinatorial structures into components. For an overview with examples see the surveys [1, 7, 13]. In this note we want to explore a correspondence between the distribution of integer divisors and the degree distribution of polynomial divisors over finite fields.

Let $\mathbb{F}_q$ be the finite field with q elements. What proportion of polynomials of degree n over $\mathbb{F}_q$ have divisors of every degree below n ? How does this question relate to the distribution of divisors of integers?

Let F be a monic polynomial of degree n over $\mathbb{F}_q$ with factorization $F = \prod_{1 \leq i \leq j} P_i$, where the P_i are irreducible monic polynomials. The set of degrees of divisors of F is given by

$$A_1 = A_1(F) = \left\{ \sum_{1 \leq i \leq j} \varepsilon_i \deg(P_i) : \varepsilon_i \in \{0, 1\} \right\} \subseteq [0, n].$$

Hence F has a divisor of every degree below n if and only if $A_1 = [0, n] \cap \mathbb{Z}$, i.e. the set A_1 has no gaps of size greater than 1.

For an integer N with prime factorization $N = \prod_{1 \leq i \leq j} p_i$, the set

$$A_2 = A_2(N) = \left\{ \sum_{1 \leq i \leq j} \varepsilon_i \log p_i : \varepsilon_i \in \{0, 1\} \right\} \subseteq [0, \log N]$$

has no gaps of size greater than 1 if and only if every interval of the form $[x, ex)$, $1 \leq x \leq N$, contains a divisor of N , i.e. the sequence of divisors of

Date: July 8, 2015.

2010 Mathematics Subject Classification. 11T06, 11N25, 05A05.

N , $1 = d_1 < d_2 < \dots < d_k = N$, satisfies

$$(1) \quad \max_{1 \leq i < k} \frac{d_{i+1}}{d_i} \leq e.$$

The similarity between the sets A_1 and A_2 , with the logarithms of the integers ($\log N$ and $\log p_i$) taking the roles of the degrees of the polynomials (n and $\deg(P_i)$), suggests that the study of polynomials having divisors of every degree is related to the study of integers whose sequence of divisors satisfies (1). Improving on earlier estimates by Tenenbaum [15, 16] and Saias [14], we found [22, Corollary 1.1] that the number of such integers $\leq x$ is given by

$$\frac{c_0 x}{\log x} \left(1 + O \left(\frac{1}{\log x} \right) \right),$$

for some positive constant c_0 . Here we use a similar strategy to establish the analogous result for polynomials.

Theorem 1. *The proportion of polynomials of degree n over $\mathbb{F}_q$, which have a divisor of every degree below n , is given by*

$$\frac{c_q}{n} \left(1 + O \left(\frac{1}{n} \right) \right).$$

The factor c_q depends only on q and satisfies

$$0 < c_q = C + O(q^{-2\tau}),$$

where $C = (1 - e^{-\gamma})^{-1} = 2.280291\dots$, γ denotes Euler's constant and the constant $\tau = 0.205466\dots$ is defined in Proposition 9.

In Theorem 1 and below, the implied constants in the error terms are absolute. In particular, all of our estimates are valid uniformly in q .

The method in [22] relies on a sharp estimate by Tenenbaum [17, Corollary 7.6, Section III.6] for the number of integers $\leq x$ which are free of prime divisors $\leq y$, for the entire range of y -values $2 \leq y \leq x$. The corresponding results for polynomials, which are available in the literature, are not precise enough for our method to succeed. Thus we first derive an improved estimate for $r(n, m)$, the proportion of polynomials of degree n over $\mathbb{F}_q$, all of whose non-constant divisors have degree $> m$. Such polynomials are sometimes referred to as m -rough, and estimates of $r(n, m)$ play a role in the analysis of factorization algorithms [6, 11]. Hence Theorem 2 may be of independent interest.

We will need the following notation. Buchstab's function $\omega(u)$ is the unique continuous solution to the equation

$$(u\omega(u))' = \omega(u-1) \quad (u > 2)$$

with initial condition $\omega(u) = 1/u$ for $1 \leq u \leq 2$. Let $\omega(u) = 0$ for $u < 1$. The number of monic irreducible polynomials of degree n over $\mathbb{F}_q$ is given

by [8, Theorem 3.25]

$$I_n = \frac{1}{n} \sum_{d|n} \mu\left(\frac{n}{d}\right) q^d \quad (n \geq 1),$$

where μ is the Möbius function. An easy exercise [8, p. 142, Ex. 3.26 and Ex. 3.27] shows that

$$(2) \quad \frac{q^n}{n} - \frac{2q^{n/2}}{n} < I_n \leq \frac{q^n}{n} \quad (n \geq 1).$$

Define

$$\lambda_q(m) := \prod_{k=1}^m \left(1 - \frac{1}{q^k}\right)^{I_k}.$$

Theorem 2. *Let $u = n/m$. For $n > m \geq 1$ we have*

$$r(n, m) = \lambda_q(m) e^\gamma \omega(u) \left(1 + O\left(\frac{(u/e)^{-u}}{m}\right)\right),$$

and

$$\lambda_q(m) = e^{-H_m} \left(1 + O\left(\frac{1}{mq^{(m+1)/2}}\right)\right) = \frac{e^{-\gamma}}{m} \left(1 + O\left(\frac{1}{m}\right)\right),$$

where $H_m := \sum_{1 \leq k \leq m} \frac{1}{m}.$

With the second estimate for $\lambda_q(m)$, Theorem 2 simplifies to

Corollary 1. *For $n > m \geq 1$ we have*

$$r(n, m) = \frac{\omega(u)}{m} \left(1 + O\left(\frac{1}{m}\right)\right).$$

Warlimont [20, Eqs. (3) and (4)] showed that Corollary 1 and the second estimate for $\lambda_q(m)$ hold in the more general setting of arithmetical semi-groups.

Inserting the estimate for $\omega(u)$ from Lemma 1 into Theorem 2, we obtain the following improvement of [6, Theorem 3.1] and [11, Theorem 3.1].

Corollary 2. *For $n > m \geq 1$ we have*

$$r(n, m) = \lambda_q(m) \left(1 + O((u/e)^{-u})\right).$$

When $m \geq 3$, we can replace $(u/e)^{-u}$ by u^{-u} in the error terms of Theorem 2 and Corollary 2.

For $m \leq n/\log n$, we derive Theorem 2 by applying the residue theorem to the generating function of $r(n, m)$. When $m > n/\log n$, we show that $r(n, m)$ is very close to $p(n, m)$, the proportion of permutations of n objects which have no cycles of length $\leq m$. The result then follows from a recent estimate of $p(n, m)$ due to Manstavičius and Petuchovas [10].

With very little extra effort, we can generalize Theorem 1 as follows. Let $f(n, m) = f_q(n, m)$ be the proportion of monic polynomials F of degree n over $\mathbb{F}_q$, with the property that the set of degrees of divisors of F (i.e. the set A_1) has no gaps of size greater than m . The original question corresponds to $m = 1$. The corresponding generalization in the case of the divisors of integers would be to replace the upper bound e in (1) by some parameter t . The asymptotic behavior of the number of integers up to x , all of whose ratios of consecutive divisors are at most t , is described in [22, Theorem 1.3] in terms of a function $d(u)$, which is defined as follows.

Let $d(u) = 0$ for $u < 0$ and

$$(3) \quad d(u) = 1 - \int_0^{\frac{u-1}{2}} \frac{d(v)}{v+1} \omega\left(\frac{u-v}{v+1}\right) dv \quad (u \geq 0),$$

where $\omega(u)$ is Buchstab's function. In [21, Theorem 1], we used equation (3) to show that

$$(4) \quad d(u) = \frac{C}{u+1} \left(1 + O(u^{-2})\right) \quad (u \geq 1),$$

where $C = (1 - e^{-\gamma})^{-1} = 2.280291\dots$, as in Theorem 1.




FIGURE 1. The graphs of $d(u)$ (solid) and $\frac{C}{u+1}$ (dashed).

The following result is the polynomial analogue of [22, Theorem 1.3].

Theorem 3. *For $n \geq 0$, $m \geq 1$ we have*

$$f(n, m) = \eta_q(m) d(n/m) \left(1 + O\left(\frac{1}{n+m}\right)\right).$$

The factor $\eta_q(m)$ depends only on q and m , and satisfies

$$0 < \eta_q(m) = 1 + O\left(\frac{1}{mq^{(m+1)\tau}}\right),$$

where $\tau = 0.205466\dots$ is defined in Proposition 9.

Depending on the size of m relative to n , we can simplify Theorem 3 in different ways. With the estimate (4), we get

Corollary 3. *For $n \geq m \geq 1$ we have*

$$f(n, m) = \frac{c_q(m)m}{n+m} \left(1 + O\left(\frac{m^2}{n^2} + \frac{1}{n}\right) \right),$$

where

$$0 < c_q(m) := C\eta_q(m) = C + O\left(\frac{1}{mq^{(m+1)\tau}}\right).$$

Theorem 1 follows from Corollary 3 with $m = 1$ and $c_q = c_q(1)$. Corollary 3 clearly implies

Corollary 4. *For $n \geq m \geq 1$ we have*

$$f(n, m) = \frac{Cm}{n+m} \left(1 + O\left(\frac{m^2}{n^2} + \frac{1}{n} + \frac{1}{mq^{(m+1)\tau}}\right) \right).$$

Finally, the estimate for $\eta_q(m)$ in Theorem 3 implies

Corollary 5. *For $n \geq 0$, $m \geq 1$ we have*

$$f(n, m) = d(n/m) \left(1 + O\left(\frac{1}{n+m} + \frac{1}{mq^{(m+1)\tau}}\right) \right).$$

We now turn to analogous results in the context of permutations and their decomposition into disjoint cycles. Let S_n be the symmetric group on $\{1, 2, \dots, n\}$. One reason for considering permutations is that our proof of the asymptotic result for the factor $\eta_q(m)$ in Theorem 3 (and hence for the factor c_q in Theorem 1) depends on Theorem 5, the corresponding result in the context of permutations. Moreover, each of the following estimates related to permutations coincides with the limit, as $q \rightarrow \infty$, of the corresponding estimate related to polynomials over $\mathbb{F}_q$. Theorem 1 takes the following form.

Theorem 4. *The proportion of permutations $\sigma \in S_n$, which have a cycle structure such that every positive integer below n can be expressed as a sum of lengths of distinct cycles of σ , is given by*

$$\frac{C}{n} \left(1 + O\left(\frac{1}{n}\right) \right),$$

where C is as in Theorem 1.

An alternate way to state Theorem 4 is as follows. The proportion of permutations $\sigma \in S_n$, with the property that for every positive integer $m \leq n$ there exists a set $M \subseteq \{1, 2, 3, \dots, n\}$ with cardinality m such that $\sigma(M) = M$, is given by $Cn^{-1} + O(n^{-2})$.

The analogue of Theorem 2 in the context of permutations is Proposition 4, which depends on a recent result by Manstavičius and Petuchovas [10].

The analogue of Theorem 3 also holds in this context. Assume a permutation $\sigma \in S_n$ has cycles of length $l_1, l_2, \dots, l_j$. The set

$$A_3 = A_3(\sigma) = \left\{ \sum_{1 \leq i \leq j} \varepsilon_i l_i : \varepsilon_i \in \{0, 1\} \right\} \subseteq [0, n]$$

represents the set of cardinalities of sets $M \subseteq \{1, 2, 3, \dots, n\}$ such that $\sigma(M) = M$. Let $g(n, m)$ denote the proportion of permutations in S_n with the property that A_3 has no gaps of size greater than m .

Theorem 5. *For $n \geq 0$, $m \geq 1$ we have*

$$g(n, m) = d(n/m) \left(1 + O\left(\frac{1}{n+m}\right) \right).$$

Unlike Theorem 3 and [22, Theorem 1.3], Theorem 5 does not involve a factor $\eta(m)$, which suggests that this variant of the problem, dealing with permutations, exhibits the simplest behavior. With the estimate (4), we get

Corollary 6. *For $n \geq m \geq 1$ we have*

$$g(n, m) = \frac{Cm}{n+m} \left(1 + O\left(\frac{m^2}{n^2} + \frac{1}{n}\right) \right).$$

Theorem 4 follows from Corollary 6 with $m = 1$.

The situation is somewhat different in the context of integer partitions. Erdős and Szalay [4], and later Dixmier and Nicolas [3], investigated the problem analogous to Theorems 1 and 4. For the integer partition $n = l_1 + l_2 + \dots + l_j$, consider the set

$$A_4 = \left\{ \sum_{1 \leq i \leq j} \varepsilon_i l_i : \varepsilon_i \in \{0, 1\} \right\} \subseteq [0, n].$$

Dixmier and Nicolas [3] call a partition *practical* if $A_4 = [0, n] \cap \mathbb{Z}$. They show that the proportion of partitions of n which are practical is given by (see [3, Theorem 2] for the full result)

$$1 - \frac{\pi}{\sqrt{6n}} + O\left(\frac{1}{n}\right).$$

Thus, unlike with integers, polynomials and permutations, almost all partitions satisfy $A_4 = [0, n] \cap \mathbb{Z}$.

A related problem is the study of the sequence of degrees n for which the particular polynomial $X^n - 1$ has divisors of every degree in $K[X]$, where K is a given field. By adapting the work of Saias [14] on practical numbers, Thompson [18] found that the number of such $n \leq x$ is $\asymp x/\log x$ when $K = \mathbb{Q}$. This was extended to any number field K by Pollack and Thompson [12]. When $K = \mathbb{F}_p$, the true order of magnitude of the number of such $n \leq x$ is still unknown, but Thompson [19] shows that they have asymptotic density zero by assuming the Generalized Riemann Hypothesis.

Acknowledgements. The author thanks Paul Pollack for suggesting the question which is answered in Theorem 1, and Eugenijus Manstavičius for providing some references.

2. PROOF OF THEOREM 2

When $m \leq n/\log n$, the first statement in Theorem 2 follows from Lemma 1 and Proposition 1. When $n/\log n < m < n$, Theorem 2 follows from Propositions 3, 6 and 7. The second statement in Theorem 2 is Proposition 7. We put $r(0, m) = 1$ for $m \geq 0$.

Lemma 1. *For $u > 1$, we have $\omega(u) - e^{-\gamma} \ll \exp(-u \log(u \log u) + O(u))$.*

Proof. This follows from [17, Theorem III.6.4]. $\square$

Proposition 1. *Let $u = n/m$. For $1 \leq m \leq n/\log n$,*

$$r(n, m) = \lambda_q(m) \left(1 + O((u/e)^{-u} m^{-1}) \right).$$

If $m \geq 3$, we can replace e by 1 in the error term.

Proof. The number of monic polynomials of degree n , which have no non-constant divisors of degree $\leq m$, is given by $[z^n]F_m(z)$, the coefficient of z^n in the power series of

$$F_m(z) := \prod_{k>m} (1 - z^k)^{-I_k} = \frac{1}{1 - qz} \prod_{1 \leq k \leq m} (1 - z^k)^{I_k} \quad (|z| < 1/q).$$

Cauchy's residue theorem yields

$$r(n, m) = \frac{[z^n]F_m(z)}{q^n} = [z^n]F_m(z/q) = \frac{1}{2\pi i} \int_{|z|=1/2} F_m(z/q) \frac{dz}{z^{n+1}}.$$

Stretching the contour to $|z| = R > 1$ leaves a residue of $\lambda_q(m)$ from the pole at $z = 1$. When $|z| = R$ we have

$$|F_m(z/q)| \leq \frac{1}{R-1} \prod_{1 \leq k \leq m} \left(1 + \frac{R^k}{q^k} \right)^{I_k} \leq \frac{1}{R-1} \exp \left(\sum_{1 \leq k \leq m} \frac{R^k}{k} \right),$$

since $I_k \leq q^k/k$ by (2) and $\log(1+x) \leq x$ for $x > 0$. Hence

$$(5) \quad |r(n, m) - \lambda_q(m)| \leq \frac{1}{R-1} \exp \left(\sum_{1 \leq k \leq m} \frac{R^k}{k} \right) \frac{1}{R^n}.$$

If $m = 1$ or 2 , the choice $R^m = n$ makes the right-hand side of (5) $\ll (u/e)^{-u} m^{-2}$. For $m \geq 3$, we write

$$\begin{aligned}
 \sum_{1 \leq k \leq m} \frac{R^k}{k} &= H_m + \sum_{1 \leq k \leq m} \frac{R^k - 1}{k} = H_m + \sum_{1 \leq k \leq m} \int_0^{\log R} e^{kt} dt \\
 &= H_m + \int_0^{\log R} (e^t - 1 + 1) \frac{e^{mt} - 1}{e^t - 1} dt \\
 &= H_m + \int_0^{\log R} (e^{mt} - 1) dt + \int_0^{\log R} \frac{e^{mt} - 1}{e^t - 1} dt \\
 &\leq H_m + \frac{R^m}{m} + \int_0^{\log R^m} \frac{e^t - 1}{t} dt,
 \end{aligned}
 \tag{6}$$

since $e^t - 1 \geq t$. A little calculus exercise shows that for $x > 0$ we have

$$\int_0^x \frac{e^t - 1}{t} dt \leq \frac{e^x}{x} \left(1 + \frac{2}{x}\right).$$

Hence

$$\sum_{1 \leq k \leq m} \frac{R^k}{k} \leq H_m + \frac{R^m}{m} + \frac{R^m}{\log R^m} \left(1 + \frac{2}{\log R^m}\right).$$

When $3 \leq m \leq \log n / \log \log \log n$, we choose R such that $R^m = n$. If $\log n / \log \log \log n < m \leq n / \log n$, we choose R such that $R^m = u \log u$. In each case, (7) shows that the right-hand side of (5) is $\ll u^{-u} m^{-2}$. $\square$

Lemma 2. For $n > m \geq 1$,

$$n r(n, m) = 1 + \sum_{m < k < n-m} r(k, m) + O\left(\frac{1}{q^{n/2}} + \frac{1}{mq^{(m+1)/2}}\right).$$

Proof. Let $R(n, m) = q^n r(n, m)$, the number of monic polynomials of degree n , which have no non-constant divisors of degree $\leq m$. As in the proof of Proposition 1, the generating function of $R(n, m)$ is given by

$$\sum_{n \geq 0} R(n, m) z^n = F_m(z) = \prod_{k > m} (1 - z^k)^{-I_k} = \exp\left(-\sum_{k > m} I_k \log(1 - z^k)\right).$$

Differentiating with respect to z gives

$$\sum_{n \geq 1} n R(n, m) z^{n-1} = F'_m(z) = F_m(z) \sum_{k > m} I_k \frac{k z^{k-1}}{1 - z^k}$$

and hence

$$\sum_{n \geq 1} n R(n, m) z^{n-1} = \left(\sum_{n \geq 0} R(n, m) z^n\right) \left(\sum_{k > m} k I_k \sum_{j \geq 1} z^{kj-1}\right).$$

Comparing coefficients of z^{n-1} , we find that

$$n R(n, m) = \sum_{k > m} k I_k \sum_{j \geq 1} R(n - kj, m).$$

We estimate $k I_k$ by (2) and divide both sides by q^n to get
(8)

$$n r(n, m) = \sum_{k > m} r(n - k, m) + O \left(\sum_{m < k \leq n} q^{-k/2} r(n - k, m) + \sum_{j \geq 2} \sum_{k > m} q^{-k(j-1)} \right).$$

Note that

$$\sum_{k > m} r(n - k, m) = 1 + \sum_{m < k < n - m} r(k, m),$$

since $r(0, m) = 1$ and $r(k, m) = 0$ for $1 \leq k \leq m$. The double sum in the error term of (8) is $\ll q^{-(m+1)}$, since $q \geq 2$. We have $r(n - k, m) = 1$ when $k = n$ and $r(n - k, m) \ll 1/m$ when $k < n$. Hence the first sum in the error term of (8) is acceptable. $\square$

Recall that $p(n, m)$ denotes the proportion of permutations of n objects having no cycles of length $\leq m$. We put $p(0, m) = 1$.

Proposition 2. *Let $u = n/m$. For $1 \leq m \leq n/\log n$,*

$$p(n, m) = e^{-H_m} \left(1 + O((u/e)^{-u} m^{-1}) \right).$$

If $m \geq 3$, we can replace e by 1 in the error term.

Proof. The generating function for $p(n, m)$ satisfies [5, Ex. IV.9]
(9)

$$D_m(z) := \sum_{n \geq 0} p(n, m) z^n = \exp \left(\sum_{k \geq m+1} \frac{z^k}{k} \right) = \frac{1}{1-z} \exp \left(- \sum_{1 \leq k \leq m} \frac{z^k}{k} \right),$$

for $|z| < 1$. Cauchy's residue theorem yields

$$p(n, m) = \frac{1}{2\pi i} \int_{|z|=1/2} D_m(z) \frac{dz}{z^{n+1}}.$$

Stretching the contour to $|z| = R > 1$ leaves a residue of e^{-H_m} from the pole at $z = 1$. When $|z| = R$ we have

$$|D(z)| \leq \frac{1}{R-1} \exp \left(\sum_{1 \leq k \leq m} \frac{R^k}{k} \right),$$

hence

$$|p(n, m) - e^{-H_m}| \leq \frac{1}{R-1} \exp \left(\sum_{1 \leq k \leq m} \frac{R^k}{k} \right) \frac{1}{R^n}.$$

The remainder of the proof is identical to that of Proposition 1 following the estimate (5). $\square$

The following result is a somewhat weaker version of a recent result by Manstavičius and Petuchovas [10, Theorem 1.1].

Proposition 3 (Manstavičius and Petuchovas). *Let $u = n/m$. For $n > m \geq \sqrt{n \log n}$ we have*

$$p(n, m) = e^{\gamma - H_m} \omega(u) \left(1 + O\left(\frac{u^{-u}}{m}\right) \right).$$

Combining Propositions 2 and 3 with Lemma 1, we get the following estimate.

Proposition 4. *Let $u = n/m$. For $n > m \geq 1$ we have*

$$p(n, m) = e^{\gamma - H_m} \omega(u) \left(1 + O\left(\frac{(u/e)^{-u}}{m}\right) \right).$$

The next estimate is an immediate consequence of Manstavičius [9, Theorem 3]. It also follows from Proposition 4.

Proposition 5 (Manstavičius). *For $n > m \geq 1$, we have*

$$p(n, m) = \frac{\omega(n/m)}{m} \left(1 + O\left(\frac{1}{m}\right) \right).$$

Lemma 3. *For $n > m \geq 0$,*

$$n p(n, m) = 1 + \sum_{m < k < n-m} p(k, m).$$

Proof. From (9) we have

$$D'_m(z) = \sum_{n \geq 1} n p(n, m) z^{n-1} = D_m(z) \sum_{k \geq m} z^k.$$

Comparing the coefficients of z^{n-1} in the last equation leads to

$$n p(n, m) = \sum_{m \leq k \leq n-1} p(n-1-k, m) = 1 + \sum_{m < k < n-m} p(k, m),$$

for $n > m \geq 0$, since $p(0, m) = 1$ and $p(k, m) = 0$ for $1 \leq k \leq m$. □

Proposition 6. *For $n \geq 1, m \geq 1$,*

$$r(n, m) = p(n, m) + O\left(\frac{1}{nq^{n/2}} + \frac{1}{m^2q^{(m+1)/2}}\right).$$

Proof. If $m \geq n \geq 1$, then $r(n, m) = p(n, m) = 0$. Hence we may assume $n > m \geq 1$. Let $s(n, m) = r(n, m) - p(n, m)$. Lemmas 2 and 3 imply

$$(10) \quad |s(n, m)| \leq \frac{B}{nq^{n/2}} + \frac{B}{nmq^{(m+1)/2}} + \frac{1}{n} \sum_{m < k < n-m} |s(k, m)|,$$

where B is the implied constant in the error term of Lemma 2. When $n/2 \leq m < n$, the last sum is empty and we have

$$(11) \quad |s(n, m)| \leq \frac{B}{nq^{n/2}} + \frac{4B}{m^2q^{(m+1)/2}}.$$

Now assume that (11) holds for all $n \geq 1$ and $n/j \leq m < n$, for some $j \geq 2$. Let $n/(j+1) \leq m < n$. Then $n \leq m(j+1)$ and $k < n - m$ imply $k < mj$. Hence (10) and the inductive hypothesis show that

$$\begin{aligned} |s(n, m)| &\leq \frac{B}{nq^{n/2}} + \frac{B}{nmq^{(m+1)/2}} + \frac{1}{n} \sum_{m < k < n-m} \left(\frac{B}{kq^{k/2}} + \frac{4B}{m^2q^{(m+1)/2}} \right) \\ &\leq \frac{B}{nq^{n/2}} + \frac{4B}{m^2q^{(m+1)/2}} \left(\frac{m}{4n} + \frac{m}{4n} \sum_{k \geq 0} q^{-k/2} + \frac{n-2m}{n} \right) \\ &\leq \frac{B}{nq^{n/2}} + \frac{4B}{m^2q^{(m+1)/2}}, \end{aligned}$$

since $\sum_{k \geq 0} q^{-k/2} \leq \sum_{k \geq 0} 2^{-k/2} < 4$. Thus (11) holds for all $n \geq 1$ and all m with $0 < m < n$. $\square$

Proposition 7. *For $m \geq 1$ we have*

$$\lambda_q(m) = e^{-H_m} \left(1 + O \left(\frac{1}{mq^{(m+1)/2}} \right) \right) = \frac{e^{-\gamma}}{m} \left(1 + O \left(\frac{1}{m} \right) \right).$$

Proof. Proposition 1 implies that $\lim_{n \rightarrow \infty} r(n, m) = \lambda_q(m)$, while Proposition 2 (or [5, Ex. IV.9]) shows that $\lim_{n \rightarrow \infty} p(n, m) = e^{-H_m}$. Letting $n \rightarrow \infty$ in Proposition 6 gives the first equation. The second equation follows from the well-known estimate $H_m = \log m + \gamma + O(m^{-1})$. $\square$

3. PROOF OF THEOREM 3

Let $\mathcal{A}(n, m)$ denote the set of monic polynomials F of degree n over $\mathbb{F}_q$ with the property that the set of degrees of divisors of F has no gaps of size greater than m . We have $f(n, m) = q^{-n} |\mathcal{A}(n, m)|$.

Lemma 4. *A monic polynomial F of degree n , $F = P_1 P_2 \cdots P_j$, where the P_i are monic irreducible polynomials with $\deg(P_1) \leq \deg(P_2) \leq \cdots \leq \deg(P_j)$, satisfies $F \in \mathcal{A}(n, m)$ if and only if*

$$(12) \quad \deg(P_i) \leq \deg(P_1 \cdots P_{i-1}) + m \quad (1 \leq i \leq j).$$

Proof. If $\deg(P_i) > \deg(P_1 \cdots P_{i-1}) + m$ for some $1 \leq i \leq j$, then the open interval $(\deg(P_1 \cdots P_{i-1}), \deg(P_i))$ is of length $> m$ and contains no degrees of divisors of F . Thus (12) is a necessary condition for $F \in \mathcal{A}(n, m)$. To see that (12) is sufficient, we use induction on j . The case $j = 1$ is obvious. Assume that (12) implies $F \in \mathcal{A}(n, m)$ for some $j \geq 1$. Let $F = P_1 P_2 \cdots P_j P_{j+1}$ satisfy (12), with j replaced by $j+1$. The set of degrees of divisors of F is $D \cup (D + \deg(P_{j+1}))$, where D is the set of degrees of divisors of $P_1 P_2 \cdots P_j$. Now D has no gaps greater than m by the inductive hypothesis and $\deg(P_{j+1}) \leq m + \max D$ by (12). Thus $D \cup (D + \deg(P_{j+1}))$ also has no gaps greater than m and $F \in \mathcal{A}(n, m)$. $\square$

Lemma 5. *For $n \geq 0$, $m \geq 0$ we have*

$$1 = \sum_{0 \leq k \leq n} f(k, m) r(n - k, k + m).$$

Proof. Lemma 4 implies that each monic polynomial F of degree n factors uniquely as $F = QR$, where $F = P_1 P_2 \cdots P_j$, the P_i are monic irreducible polynomials with $\deg(P_1) \leq \deg(P_2) \leq \dots \leq \deg(P_j)$,

$$Q = P_1 \cdots P_{i_0}, \quad R = P_{i_0+1} \cdots P_j,$$

and i_0 is the unique index such that

$$Q \in \mathcal{A}(\deg(Q), m), \quad \deg P_{i_0+1} > \deg(Q) + m.$$

The case $i_0 = 0$ corresponds to $Q = 1$ and $R = F$, while $i_0 = j$ means $Q = F \in \mathcal{A}(n, m)$ and $R = 1$. Given $\deg(Q) = k$, there are $q^k f(k, m)$ choices for Q and $q^{n-k} r(n - k, k + m)$ choices for R . Counting all of the q^n monic polynomials of degree n according to the degree of Q , we find that

$$q^n = \sum_{0 \leq k \leq n} q^k f(k, m) q^{n-k} r(n - k, k + m),$$

from which the result follows. $\square$

Lemma 6. *For $n \geq 0$, $m \geq 0$ we have*

$$1 = f(n, m) + \sum_{0 \leq k < \frac{n-m}{2}} f(k, m) r(n - k, k + m).$$

Proof. This follows from Lemma 5 since $r(0, n+m) = 1$ and $r(n-k, k+m) = 0$ if $\frac{n-m}{2} \leq k < n$. $\square$

Lemma 7. *For $m \geq 0$ we have*

$$1 = \sum_{k \geq 0} f(k, m) \lambda_q(k + m).$$

Proof. If $m = 0$, we have $f(0, 0) = 1$, $\lambda_q(0) = 1$ and $f(k, 0) = 0$ for $k \geq 1$. Now fix any $m \geq 1$. We have $r(n-k, k+m) \asymp (k+m)^{-1}$ for $n-k > k+m$, by Corollary 1. Lemma 6 implies that $\sum_{0 \leq k < \frac{n-m}{2}} \frac{f(k, m)}{k+m}$ is bounded from above as $n \rightarrow \infty$ and hence $\sum_{k \geq 0} \frac{f(k, m)}{k+m}$ converges. Thus

$$\liminf_{k \rightarrow \infty} f(k, m) = 0, \quad \lim_{n \rightarrow \infty} \sum_{n/\log n < k < \frac{n-m}{2}} f(k, m) r(n - k, k + m) = 0.$$

Lemma 6 shows that, as $n \rightarrow \infty$,

$$\begin{aligned} 1 &= o(1) + f(n, m) + \sum_{k \leq n/\log n} f(k, m) r(n - k, k + m) \\ &= o(1) + f(n, m) + \sum_{k \leq n/\log n} f(k, m) \lambda_q(k + m), \end{aligned}$$

by Corollary 2. Since $\liminf_{n \rightarrow \infty} f(n, m) = 0$ and the last sum is increasing in n , the result follows. $\square$

For real $t \geq 0$, we define $f(t, m) = f(\lfloor t \rfloor, m)$.

Lemma 8. *For $n \geq 0$, $m \geq 1$ we have*

$$f(n, m) = \int_0^\infty \frac{f(t, m)}{t + m} \left(e^{-\gamma} - \omega \left(\frac{n + m}{t + m} - 1 \right) \right) dt + E(n, m),$$

where

$$E(n, m) \ll E_0(n, m) := \int_0^\infty \frac{f(t, m)}{(t + m)^2} \exp \left(-\frac{n + m}{t + m} \right) dt + \frac{f(\frac{n-m}{2}, m)}{n + m}.$$

Proof. Lemmas 6 and 7 imply

$$\begin{aligned} f(n, m) &= \sum_{0 \leq k < \frac{n-m}{2}} f(k, m) \left(\lambda_q(k + m) - r(n - k, k + m) \right) \\ &\quad + \sum_{k \geq \frac{n-m}{2}} f(k, m) \lambda_q(k + m) =: S_1 + S_2, \end{aligned}$$

say. We approximate $r(n - k, k + m)$ in S_1 by Theorem 2 to get

$$S_1 = \sum_{0 \leq k < \frac{n-m}{2}} f(k, m) \lambda_q(k + m) e^\gamma \left(e^{-\gamma} - \omega \left(\frac{n - k}{k + m} \right) \right) + E_1(n, m),$$

where $E_1(n, m) \ll E_0(n, m)$. Next, we approximate $\lambda_q(k + m)$ in S_1 and S_2 by the second estimate in Proposition 7. The resulting error term is again $\ll E_0(n, m)$ by Lemma 1. Thus

$$f(n, m) = S_1 + S_2 = \sum_{k \geq 0} \frac{f(k, m)}{k + m} \left(e^{-\gamma} - \omega \left(\frac{n + m}{k + m} - 1 \right) \right) + E_2(n, m),$$

where $E_2(n, m) \ll E_0(n, m)$, since $\omega(1) = 1$ and $\omega(u) = 0$ for $u < 1$. It remains to replace the sum by an integral. If $g(k)$ represents the k -th term of the sum, we estimate the error simply by $g(k) - \int_k^{k+1} g(t) dt \ll \max_{t \in [k, k+1]} |g'(t)|$ and use Lemma 1 and $\omega'(u) \ll e^{-2u}$ for $u \geq 1$ (see [17, Theorem III.6.4]). The discontinuity of $\omega(u)$ at $u = 1$ leads to an error of $\ll f(\frac{n-m}{2}, m)/(n + m)$. Thus we find that the error from replacing the sum by the integral is also $\ll E_0(n, m)$. $\square$

Proof of Theorem 3. Note that replacing the integer n in Lemma 8 by the real number x with $n = \lfloor x \rfloor$ leaves the left-hand-side unchanged, and alters the integral by $\ll E_0(x, m)$. Thus, for $x \geq 0$,

$$f(x, m) = \alpha(m) - \int_0^\infty \frac{f(t, m)}{t + m} \omega \left(\frac{x + m}{t + m} - 1 \right) dt + E(x, m),$$

where

$$\alpha(m) := e^{-\gamma} \int_0^\infty \frac{f(t, m)}{t + m} dt$$

and $E(x, m) \ll E_0(x, m)$. Lemma 7 ensures that the integral defining $\alpha(m)$ converges and that $\alpha(m) \ll 1$. Let $z \geq 0$ be given by $e^z = x/m + 1$ and $y \geq 0$ by $e^y = t/m + 1$. Define $G_m(z) := f(m(e^z - 1), m) e^z$ and $\Omega(z) := \omega(e^z - 1)$. We get

$$G_m(z) = \alpha(m)e^z - \int_0^z G_m(y) \Omega(z - y) e^{z-y} dy + E_m(z),$$

where

$$(13) \quad E_m(z) \ll \frac{e^z}{m} \int_0^\infty \frac{G_m(y)}{e^{2y}} \exp(-e^{z-y}) dy + \frac{G_m(z - \log 2)}{me^z}.$$

The last equation allows us to calculate Laplace transforms:

$$\widehat{G}_m(s) = \frac{\alpha(m)}{s-1} - \widehat{G}_m(s) \widehat{\Omega}(s-1) + \widehat{E}_m(s) \quad (\operatorname{Re} s > 1).$$

Hence

$$\widehat{G}_m(s) = \frac{\alpha(m)}{(s-1)(1 + \widehat{\Omega}(s-1))} + \frac{\widehat{E}_m(s)}{1 + \widehat{\Omega}(s-1)} \quad (\operatorname{Re} s > 1).$$

Equation (3) written in terms of $G(z) := e^z d(e^z - 1) \asymp 1$ is

$$G(z) = e^z - \int_0^z G(y) \Omega(z - y) e^{z-y} dy,$$

which shows that the Laplace transform of $G(z)$ is given by

$$\widehat{G}(s) = \frac{1}{(s-1)(1 + \widehat{\Omega}(s-1))} \quad (\operatorname{Re} s > 1).$$

Thus

$$\begin{aligned} \widehat{G}_m(s) &= \alpha(m) \widehat{G}(s) + \widehat{E}_m(s) \widehat{G}(s)(s-1) \\ &= \alpha(m) \widehat{G}(s) + \widehat{E}_m(s) (\widehat{G}'(s) - \widehat{G}(s) + 1), \end{aligned}$$

since $G(0) = 1$. Now

$$G'(y) - G(y) = e^{2y} d'(e^y - 1) = -C + O(e^{-2y})$$

by [21, Corollary 5]. Inversion of the Laplace transforms yields

$$(14) \quad G_m(z) = \alpha(m)G(z) + E_m(z) + \int_0^z E_m(y) \left(-C + O(e^{-2(z-y)}) \right) dy.$$

Since $0 \leq f(x, m) \leq 1$, we have $0 \leq G_m(z) \leq e^z$. Thus (13) shows that $E_m(z) \ll 1$ and (14) yields $G_m(z) \ll 1 + z$. Using this estimate in (13) we find that $E_m(z) \ll (1+z)e^{-z}$ and (14) now gives $G_m(z) \ll 1$. Appealing to (13) one last time, we finally arrive at

$$E_m(z) \ll \frac{1}{me^z}.$$

Hence

$$\beta(m) := - \int_0^\infty E_m(y) dy = - \int_0^z E_m(y) dy + O\left(\frac{1}{me^z}\right)$$

and

$$\int_0^z |E_m(y)| e^{-2(z-y)} dy \ll \frac{1}{me^z}.$$

Thus (14) yields

$$G_m(z) = \alpha(m)G(z) + C\beta(m) + O\left(\frac{1}{me^z}\right),$$

that is

$$f(x, m) = \alpha(m) d(x/m) + \frac{C\beta(m)}{x/m + 1} + O\left(\frac{m}{(x+m)^2}\right),$$

for $x \geq 0$, $m \geq 1$. Letting $x = m$ in the last equation shows that $\alpha(m) = 1 + O(1/m)$, since $\beta(m) \ll 1/m$, $f(m, m) = 1$ and $d(1) = 1$. With

$$(15) \quad \eta_q(m) := \alpha(m) + \beta(m) = 1 + O(1/m),$$

the estimate (4) yields

$$(16) \quad f(x, m) = \eta_q(m) d(x/m) + O\left(\frac{m}{(x+m)^2}\right).$$

We need to show that $\eta_q(m) > 0$. If $\eta_q(m) = 0$ for some $m \geq 1$ and some $q \geq 2$, then there exists an m such that $\eta_q(m) = 0$ but $\eta_q(m+1) > 0$, since $\eta_q(m) = 1 + O(1/m)$. Hence $f(n, m) \ll_{m,q} n^{-2}$ while $f(n, m+1) \asymp_{m,q} n^{-1}$. Lemma 9 shows that this is impossible. Finally, the estimate $\eta_q(m) = 1 + O(m^{-1}q^{-(m+1)\tau})$ is the topic of Proposition 9. $\square$

Lemma 9. *For $n \geq 0$, $m \geq 1$ we have $f(n, m+1) \leq qf(n+1, m)$.*

Proof. For each $F \in \mathcal{A}(n, m+1)$, we have $xF(x) \in \mathcal{A}(n+1, m)$. Hence $|\mathcal{A}(n, m+1)| \leq |\mathcal{A}(n+1, m)|$, that is $q^n f(n, m+1) \leq q^{n+1} f(n+1, m)$. $\square$

4. PROOF OF THEOREM 5

Let $\mathcal{B}(n, m)$ denote the set of permutations $\sigma \in S_n$ with the property that A_3 has no gaps of size greater than m . We have $g(n, m) = |\mathcal{B}(n, m)|/n!$. The following five lemmas correspond to Lemmas 4 through 8 in the last section. We omit some of the proofs of these lemmas and the first half of the proof of Theorem 5, since they are almost identical to those of the previous section.

Lemma 10. *A permutations $\sigma \in S_n$, which decomposes into j cycles with cycle lengths $l_1 \leq l_2 \leq \dots \leq l_j$, satisfies $\sigma \in \mathcal{B}(n, m)$ if and only if*

$$l_i \leq m + \sum_{1 \leq k < i} l_k \quad (1 \leq i \leq j).$$

Proof. Follow the proof of Lemma 4, replacing $\deg(P_i)$ by l_i . $\square$

Lemma 11. *For $n \geq 0$, $m \geq 0$ we have*

$$1 = \sum_{0 \leq k \leq n} g(k, m) p(n-k, k+m).$$

Proof. Lemma 10 implies that each $\sigma \in S_n$ decomposes uniquely as $\sigma = \rho\tau$, where $\sigma = \sigma_1\sigma_2 \cdots \sigma_j$, the σ_i are cycles with lengths $l_1 \leq l_2 \leq \dots \leq l_j$,

$$\rho = \sigma_1 \cdots \sigma_{i_0}, \quad \tau = \sigma_{i_0+1} \cdots \sigma_j,$$

and i_0 is the unique index such that

$$l_i \leq m + \sum_{1 \leq r < i} l_r \quad (1 \leq i \leq i_0), \quad l_{i_0+1} > l_1 + \dots + l_{i_0} + m.$$

The case $i_0 = 0$ corresponds to $\tau = \sigma$, while $i_0 = j$ means $\rho = \sigma \in \mathcal{B}(n, m)$. Given that $\sum_{1 \leq r \leq i_0} l_r = k$, there are $\binom{n}{k}$ ways to choose the k numbers from $\{1, 2, \dots, n\}$, which ρ is acting on. Once these k numbers are chosen, there are $k!g(k, m)$ choices for ρ and $(n-k)!p(n-k, k+m)$ choices for τ . Counting all of the $n!$ permutations in S_n according to $k = \sum_{1 \leq r \leq i_0} l_r$, we find that

$$n! = \sum_{0 \leq k \leq n} \binom{n}{k} k! g(k, m) (n-k)! p(n-k, k+m),$$

from which the result follows. $\square$

Lemma 12. *For $n \geq 0$, $m \geq 0$ we have*

$$1 = g(n, m) + \sum_{0 \leq k < \frac{n-m}{2}} g(k, m) p(n-k, k+m).$$

Proof. This follows from Lemma 11 since $p(0, n+m) = 1$ and $p(n-k, k+m) = 0$ if $\frac{n-m}{2} \leq k < n$. $\square$

Lemma 13. *For $m \geq 0$ we have*

$$1 = \sum_{k \geq 0} g(k, m) e^{-H_{k+m}}.$$

Proof. Follow the proof of Lemma 7. $\square$

Lemma 14. *For $n \geq 0$, $m \geq 1$ we have*

$$g(n, m) = \int_0^\infty \frac{g(t, m)}{t+m} \left(e^{-\gamma} - \omega \left(\frac{n+m}{t+m} - 1 \right) \right) dt + \mathcal{E}(n, m),$$

where

$$\mathcal{E}(n, m) \ll \mathcal{E}_0(n, m) := \int_0^\infty \frac{g(t, m)}{(t+m)^2} \exp \left(-\frac{n+m}{t+m} \right) dt + \frac{g(\frac{n-m}{2}, m)}{n+m}.$$

Proof. Follow the proof of Lemma 8 and use Proposition 4 (in place of Theorem 2) to approximate $p(n-k, k+m)$. $\square$

Proof of Theorem 5. As in the proof of Theorem 3, we arrive at the analogue of (15) and (16), namely

$$(17) \quad g(x, m) = \eta(m) d(x/m) + O \left(\frac{m}{(x+m)^2} \right),$$

where $\eta(m) = 1 + O(1/m)$. It remains to show that $\eta(m) = 1$ for all $m \geq 1$. To this end, we multiply the equation in Lemma 11 by z^n , $|z| < 1$, to get

$$z^n = \sum_{0 \leq k \leq n} g(k, m) z^k p(n - k, k + m) z^{n-k}.$$

Summing over $n \geq 0$ yields

$$\frac{1}{1 - z} = \sum_{k \geq 0} g(k, m) z^k \frac{1}{1 - z} \exp \left(- \sum_{j=1}^{k+m} \frac{z^j}{j} \right),$$

by (9), and hence

$$z^m = \sum_{k \geq m} g(k - m, m) z^k \exp \left(- \sum_{j=1}^k \frac{z^j}{j} \right).$$

Differentiating both sides with respect to z , we obtain

$$(18) \quad m z^{m-1} = \sum_{k \geq m} g(k - m, m) z^k \exp \left(- \sum_{j=1}^k \frac{z^j}{j} \right) \left(\frac{k}{z} - \frac{1 - z^k}{1 - z} \right).$$

Our plan is to learn about $\eta(m)$ by inserting the estimate (17) into (18). Let $m \geq 1$ be arbitrary but fixed and let $z = 1 - \delta$, where $0 < \delta < 1/2$. We will need the following estimates. First,

$$\frac{k}{z} - \frac{1 - z^k}{1 - z} = \frac{k}{1 - \delta} - \frac{1 - \exp(-k\delta(1 + O(\delta)))}{\delta} \ll \begin{cases} k^2\delta & (k \leq \delta^{-1}) \\ k & (k > \delta^{-1}). \end{cases}$$

Second, as in (6), we have

$$\begin{aligned} \sum_{j=1}^k \frac{z^j}{j} &= H_k + \int_{\log z}^0 (1 - e^{kt}) dt - \int_{\log z}^0 \frac{1 - e^{kt}}{1 - e^t} dt \\ &\geq H_k + 0 - \int_{\log z}^0 \frac{1 - e^{kt}}{-t} (1 + O(t)) dt \\ &\geq \log k - \max(0, \log(-k \log z)) - O(1) \\ &\geq \min(\log \delta^{-1}, \log k) - O(1), \end{aligned}$$

hence

$$\exp \left(- \sum_{j=1}^k \frac{z^j}{j} \right) \ll \max(\delta, 1/k).$$

Since $g(k - m, m) \ll m/k$ by (17), the contribution from $k \leq \delta^{-1/2}$ to the right-hand side of (18) is

$$\ll \sum_{m \leq k \leq \delta^{-1/2}} \frac{m}{k} \cdot 1 \cdot \frac{1}{k} \cdot k^2 \delta \leq m \delta^{1/2}.$$

For $k > \delta^{-1/2}$, we estimate $g(k - m, m)$ in (18) by

$$g(k - m, m) = \frac{C\eta(m)m}{k} + O\left(\frac{m}{k^2} + \frac{m^3}{k^3}\right),$$

which follows from (17) and (4). The contribution from the error terms is

$$\ll \sum_{\delta^{-1/2} < k \leq \delta^{-1}} \frac{m^3}{k^2} \cdot 1 \cdot \frac{1}{k} \cdot k^2 \delta + \sum_{k > \delta^{-1}} \frac{m^3}{k^2} \cdot e^{-\delta k} \cdot \delta \cdot k \ll m^3 \delta \log(\delta^{-1}).$$

After inserting the terms corresponding to $1 \leq k \leq \delta^{-1/2}$, which contribute $\ll m\delta^{1/2}$, we get

$$mz^{m-1} = O\left(m^3 \delta^{1/2}\right) + \sum_{k \geq 1} \frac{C\eta(m)m}{k} z^k \exp\left(-\sum_{j=1}^k \frac{z^j}{j}\right) \left(\frac{k}{z} - \frac{1 - z^k}{1 - z}\right),$$

where $z = 1 - \delta$. Taking the limit as $\delta \rightarrow 0^+$, we find that

$$\frac{1}{C\eta(m)} = \lim_{z \rightarrow 1^-} \sum_{k \geq 1} \frac{z^k}{k} \exp\left(-\sum_{j=1}^k \frac{z^j}{j}\right) \left(\frac{k}{z} - \frac{1 - z^k}{1 - z}\right).$$

Since the right-hand side is independent of m , so is $\eta(m)$. But $\eta(m) = 1 + O(1/m)$. Thus $\eta(m) = 1$ for all $m \geq 1$ and the proof of Theorem 5 is complete. $\square$

5. AN ESTIMATE FOR $\eta_q(m)$.

Define κ to be the unique positive constant satisfying

$$(19) \quad 1 = \int_1^\infty \omega(y) (y+1)^{-1-\kappa} dy.$$

Numerical calculations show that $\kappa = 0.433489\dots$. This estimate was obtained using exact formulas for $\omega(y)$ for $0 \leq y \leq 5$, derived with the help of Mathematica; we used a table of zeros and relative extrema of $\omega(u) - e^{-\gamma}$ on the interval $[5, 10.3355]$ due to Cheer and Goldston [2], and the estimate $|\omega(u) - e^{-\gamma}| < 1/\Gamma(u+1)$ from [21, Lemma 1] for $u \geq 10.3355$.

Proposition 8. *For $n \geq 0$, $m \geq 1$ we have*

$$f(n, m) = g(n, m) + O\left(\frac{n^\kappa}{m^{1+\kappa} q^{(m+1)/2}}\right).$$

Proof. We show the result for all $m \geq 1$ and $n \leq m2^j$ by induction on j . If $n \leq m$, then $f(n, m) = g(n, m) = 1$. Assume that, for some $j \geq 0$, there is a constant $B_j \geq 1$, such that for all $m \geq 1$ and $n \leq m2^j$,

$$|f(n, m) - g(n, m)| \leq \frac{B_j(n+m)^\kappa}{m^{1+\kappa} q^{(m+1)/2}}.$$

Now let $m \geq 1$ and $m2^j < n \leq m2^{j+1}$. From Lemmas 6 and 12 we have

$$\begin{aligned} |f(n, m) - g(n, m)| &\leq \sum_{0 \leq k < \frac{n-m}{2}} \left| g(k, m) p(n-k, k+m) - f(k, m) r(n-k, k+m) \right| \\ &\leq S_1 + S_2, \end{aligned}$$

where, by Proposition 5,

$$\begin{aligned} S_1 &:= \sum_{0 \leq k < \frac{n-m}{2}} p(n-k, k+m) \left| g(k, m) - f(k, m) \right| \\ &\leq \sum_{0 \leq k < \frac{n-m}{2}} \left(\frac{\omega\left(\frac{n-k}{k+m}\right)}{k+m} + \frac{O(1)}{(k+m)^2} \right) \frac{B_j(k+m)^\kappa}{m^{1+\kappa}q^{(m+1)/2}} \\ &\leq \frac{B_j}{m^{1+\kappa}q^{(m+1)/2}} \left(\int_0^{\frac{n-m}{2}} \frac{\omega\left(\frac{n-t}{t+m}\right)}{(t+m)^{1-\kappa}} dt + O\left(\frac{1}{m^{1-\kappa}}\right) \right) \\ &\leq \frac{B_j(n+m)^\kappa}{m^{1+\kappa}q^{(m+1)/2}} \left(\int_1^\infty \omega(y) (y+1)^{-1-\kappa} dy + O\left(\frac{m^\kappa}{m(n+m)^\kappa}\right) \right) \\ &= \frac{B_j(n+m)^\kappa}{m^{1+\kappa}q^{(m+1)/2}} \left(1 + O\left(\frac{1}{m2^{j\kappa}}\right) \right). \end{aligned}$$

By Proposition 6,

$$\begin{aligned} S_2 &:= \sum_{0 \leq k < \frac{n-m}{2}} f(k, m) \left| p(n-k, k+m) - r(n-k, k+m) \right| \\ &\ll \sum_{0 \leq k < \frac{n-m}{2}} \frac{1}{(k+m)q^{(k+m+1)/2}} \ll \frac{1}{mq^{(m+1)/2}} \\ &\leq \frac{(n+m)^\kappa}{m^{1+\kappa}q^{(m+1)/2}} \frac{1}{2^{j\kappa}}. \end{aligned}$$

Combining these estimates, we obtain

$$|f(n, m) - g(n, m)| \leq \frac{B_{j+1}(n+m)^\kappa}{m^{1+\kappa}q^{(m+1)/2}} \quad (m \geq 1, n \leq m2^{j+1}),$$

where $B_{j+1} := B_j(1 + A2^{-j\kappa})$ and A is a suitable absolute constant. $\square$

Proposition 9. Let $\tau = 1/(4+2\kappa) = 0.205466\dots$, where κ is given by (19). For $m \geq 1$, we have

$$\eta_q(m) = 1 + O\left(\frac{1}{mq^{(m+1)\tau}}\right).$$

Proof. Assume $n \geq m$. In Proposition 8, replace $f(n, m)$ and $g(n, m)$ by their respective estimates (16) and (17), with $\eta(m) = 1$. We obtain

$$\eta_q(m)d(n/m) = d(n/m) + O\left(\frac{m}{n^2} + \frac{n^\kappa}{m^{1+\kappa}q^{(m+1)/2}}\right).$$

Dividing by $d(n/m) \asymp m/n$ gives

$$\eta_q(m) = 1 + O\left(\frac{1}{n} + \frac{n^{1+\kappa}}{m^{2+\kappa}q^{(m+1)/2}}\right).$$

The result now follows with $n = \lfloor mq^{(m+1)/(4+2\kappa)} \rfloor = \lfloor mq^{(m+1)\tau} \rfloor$. $\square$

REFERENCES

- [1] R. Arratia, A.D. Barbour and S. Tavaré, Random combinatorial structures and prime factorizations, *Notices Amer. Math. Soc.* **44** (1997), 903–910.
- [2] A. Y. Cheer and D. A. Goldston, A differential delay equation arising from the sieve of Eratosthenes, *Math. Comp.* **55** (1990), 129–141.
- [3] J. Dixmier and J.-L. Nicolas, Partitions without small parts. Number theory, Vol. I (Budapest, 1987), 9–33, *Colloq. Math. Soc. János Bolyai*, 51, North-Holland, Amsterdam, 1990.
- [4] P. Erdős and M. Szalay, On some problems of J. Dénes and P. Turán, *Studies in pure mathematics*, 187–212, Birkhäuser, Basel, 1983.
- [5] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge Univ. Press, 2009.
- [6] S. Gao and D. Panario, Tests and constructions of irreducible polynomials over finite fields, *Foundations of computational mathematics (Rio de Janeiro, 1997)*, 346–361, Springer, Berlin, 1997.
- [7] A. Granville, The anatomy of integers and permutations, <http://www.dms.umontreal.ca/~andrew/PDF/Anatomy.pdf>
- [8] R. Lidl and H. Niederreiter, *Finite fields*, vol. 20 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 1997.
- [9] E. Manstavičius, On permutations missing short cycles, *Lietuvos matem. rink.*, spec. issue, **42** (2002), 1–6.
- [10] E. Manstavičius and R. Petuchovas, Local probabilities and total variation distance for random permutations, preprint.
- [11] D. Panario and B. Richmond, Analysis of Ben-Or’s polynomial irreducibility test, *Random Structures Algorithms* **13** (1998), 439–456.
- [12] P. Pollack and L. Thompson, On the degrees of divisors of $T^n - 1$, *New York J. Math.* **19** (2013), 91–116.
- [13] Z. Rudnick, Some problems in analytic number theory for polynomials over a finite field, arXiv:1501.01769.
- [14] E. Saias, Entiers à diviseurs denses 1, *J. Number Theory* **62** (1997), 163–191.
- [15] G. Tenenbaum, Sur un problème de crible et ses applications, *Ann. Sci. École Norm. Sup. (4)* **19** (1986), 1–30.
- [16] G. Tenenbaum, Sur un problème de crible et ses applications, 2. Corrigendum et étude du graphe divisoriel. *Ann. Sci. École Norm. Sup. (4)* **28** (1995), 115–127.
- [17] G. Tenenbaum, *Introduction to Analytic and Probabilistic Number Theory*, Cambridge Studies in Advanced Mathematics, Vol. 46, Cambridge Univ. Press, 1995.
- [18] L. Thompson, Polynomials with divisors of every degree, *J. Number Theory* **132** (2012), 1038–1053.
- [19] L. Thompson, On the divisors of $x^n - 1$ in $F_p[x]$, *Int. J. Number Theory* **9** (2013), 421–430.

- [20] R. Warlimont, Arithmetical semigroups II: sieving by large and small prime elements. Sets of multiples, *Manuscripta Math.* **71** (1991), 197–221.
- [21] A. Weingartner, Integers with dense divisors 3, *J. Number Theory* **142** (2014), 211–222.
- [22] A. Weingartner, Practical numbers and the distribution of divisors, *Q. J. Math.* **66** (2015), 743–758.

DEPARTMENT OF MATHEMATICS, 351 WEST UNIVERSITY BOULEVARD, SOUTHERN
UTAH UNIVERSITY, CEDAR CITY, UTAH 84720, USA
E-mail address: `weingartner@suu.edu`